

THE CASE FOR $\Omega_M = 0.33 \pm 0.035$

MICHAEL S. TURNER

Departments of Astronomy & Astrophysics and of Physics, Enrico Fermi Institute
 and Center for Cosmological Physics, The University of Chicago, Chicago, IL 60637-1433
 AND

NASA/Fermilab Astrophysics Center
 Fermi National Accelerator Laboratory, Batavia, IL 60510-0500
 mturner@oddjob.uchicago.edu
Draft version February 1, 2008

ABSTRACT

For decades, the determination of the mean density of matter (Ω_M) has been tied to the distribution of light. This has led to a “bias,” perhaps as large as a factor of 2, in determining a key cosmological parameter. Recent measurements of the physical properties of clusters, cosmic microwave background (CMB) anisotropy and the power spectrum of mass inhomogeneity now allow a determination of Ω_M without “visual bias.” The early data lead to a consistent picture of the matter and baryon densities, with $\Omega_B = 0.039 \pm 0.0075$ and $\Omega_M = 0.33 \pm 0.035$.

Subject headings: cosmology: cosmological parameters, early universe

1. INTRODUCTION

The mean mass density of the Universe is a cosmological parameter of great importance. It can be expressed as a fraction of the critical density, $\Omega_M \equiv \rho_M / \rho_{\text{crit}}$, or in physical units (g cm^{-3}). Moving between the Hubble constant:

$$\rho_M = 1.88(\Omega_M h^2) \times 10^{-29} \text{ g cm}^{-3} \quad (1)$$

where as usual, $H_0 = 100h \text{ km sec}^{-1} \text{ Mpc}^{-1}$ and $\rho_{\text{crit}} = 3H_0^2/8\pi G = 1.88h^2 \times 10^{-29} \text{ g cm}^{-3} \approx 10^{-29} \text{ g cm}^{-3}$.

Over the past thirty years much effort has been devoted to determining the mean matter density (see e.g., Faber & Gallagher, 1979; Dekel et al, 1997; Turner, 2000; Primack, 2001). The task is daunting: the density of matter in a large enough sample of the Universe to be representative ($> 10^6 \text{ Mpc}^3$) must be determined. The most well developed techniques have been tied in one way or another to the distribution of light; e.g., mass-to-light ratios and peculiar velocities (see e.g., Bahcall et al, 1995; Carlberg et al, 1997; Dekel, 1994; Willick & Strauss, 1998).

The mass-to-ratio technique begins from the deceptively simple equation that relates the mean luminosity density and the mean mass density:

$$\rho_M = \langle M/L \rangle \mathcal{L}. \quad (2)$$

Using the canonical value for the B-band luminosity density today¹ and solar units, Eq. 2 can be written as the familiar,

$$\Omega_M = \langle M/L \rangle_B / 1200h. \quad (3)$$

The task is now transformed to determining the mean mass-to-light ratio. Either $\langle M/L \rangle$ must be measured in a large enough volume to be representative ($> 10^6 \text{ Mpc}^3$), or measured for a system that can be argued to be “typical.” Needless to say, the accuracy of the inferred mean matter density can be no better than that of the mean mass-to-light ratio.

In attempting to determine $\langle M/L \rangle$ attention has been focused on clusters, because of their size and relatively well

determined total masses. However, clusters are not large enough to provide a representative sample of the Universe, and because the cluster environment is a high density one (10^2 – 10^3 times the mean density) in which only a few percent of galaxies find themselves, there is no *a priori* reason to believe that their mass-to-light ratio is representative.

The CNOC sample provides the largest and best understood sample of clusters, and from it Carlberg et al (1997) infer $\Omega_M = 0.19 \pm 0.04$. By comparing cluster data with N-body simulations Bahcall et al (2000) find a similar value, $\Omega_M = 0.16 \pm 0.05$.

Field galaxies are more representative, but problematic because galaxy halos are large. In fact, no definitive evidence yet exists that the total mass of even a single galaxy halo has been determined. The advent of weak-gravitational lensing to probe the far reaches of galaxy halos and to determine their total masses may soon surmount this hurdle (see e.g., Fischer et al, 2000). However, for now, the values for Ω_M inferred from clusters (quoted above) represent the state of this art in mass-to-light ratios.

Peculiar velocities and bulk flows are directly tied to the underlying mass distribution; comparison of peculiar velocities and mass inhomogeneity ($\delta\rho/\rho$) fixes the mean density (see e.g., Dekel 1994). However, it is the distribution of galaxies – not mass – that is measured in redshift surveys. For this reason, this technique probes $\beta = \Omega_M^{0.6}/b$, where b is the assumed linear bias factor between light and mass. (Nonlinear effects break the degeneracy, but there has been little success in exploiting this fact.) The peculiar-velocity technique has its own convergence problem: Unless the mass distribution is surveyed to a volume sufficiently large to include all of the inhomogeneity that gives rise to the peculiar velocity, an accurate value for Ω_M will not be obtained. Currently, there are unexplained differences in the values obtained for β , which range from 0.4 to almost 1.

¹ Important questions still remain about the mean luminosity density, including evolution.

A first step toward a physically based technique was proposed by White et al (1993). The method goes like this. Assume that galaxy clusters provide a “fair sample” of matter in the Universe, i.e., that their composition reflects the universal baryon-to-total mass ratio ($= \Omega_B/\Omega_M$). Then, from our knowledge of the baryon density from big-bang nucleosynthesis (BBN), the mean matter density can be inferred:

$$\Omega_M = \left(\frac{M_T}{M_B} \right)_{\text{cluster}} \Omega_B(\text{BBN}) \quad (4)$$

While the bias factor b never enters, there are potential systematics: Can one find all the cluster baryons? Can the ratio be determined on large enough scales to ensure that baryon settling is not an issue? White et al (1993) and others (e.g., Mohr et al, 1998) have argued that the answer to these questions is yes, and several recent reviews of the mean matter density have given significant weight to this technique (see e.g., Dekel et al, 1997; Turner, 2000; and Primack, 2001).

Today, other physical measurements, including CMB anisotropy and measurements of the power spectrum of matter inhomogeneity, can be added to the mix. As I will describe, the physically based methods have become mature enough to self-consistently and robustly pin down the mass density with a relatively small error bar. The value I derive from the current data is:

$$\Omega_M = 0.33 \pm 0.035 \quad (1 \sigma) \quad (5)$$

There are however several potential sources of systematic error (see below). Nonetheless, I believe that this marks an important first step toward an accurate determination of the mean matter density that is independent of the relationship of mass to light.

2. THE INPUT DATA: PHYSICAL MEASUREMENTS

A suite of physical measurements can now probe the distribution of matter and the mean matter in an unbiased way. They include accurate determinations of CMB anisotropy on angular scales down to 0.1 degrees and measurements of the shape of the power spectrum of matter inhomogeneity. CMB anisotropy depends significantly upon the physical baryon and matter density, as well as other parameters, but not upon the connection between mass and light.

The shape of the power spectrum of mass inhomogeneity depends significantly upon $\Omega_M h$ and Ω_M/Ω_B ; it can be measured by weak-gravitational lensing (cosmic shear) and through the distribution of galaxies (redshift surveys). Cosmic shear measurements are insensitive to bias, but are much less mature. Redshift surveys are sensitive to bias in determining the power spectrum itself. However, the shape of the power spectrum is only sensitive to strong scale-dependent bias on large scales around the bend in the power spectrum associated with matter-radiation equality ($\sim 30 \text{ Mpc}/\Omega_M h^2$).

With MAP, Planck, full results from the SDSS and 2dF, and additional cosmic-shear measurements coming, the future for an accurate, physically based measurement of Ω_M is very bright. I believe there is now enough data to make a preliminary estimate of Ω_M using these techniques.

The following physical measurements comprise my input data for determining the baryon and total matter den-

ties:

Power Spectrum	$\Omega_M h = 0.20 \pm 0.03$
	$\Omega_B/\Omega_M = 0.15 \pm 0.07$
CMB Anisotropy	$\Omega_M h^2 = 0.16 \pm 0.04$
	$\Omega_B h^2 = 0.022^{+0.004}_{-0.003}$
BBN	$\Omega_B h^2 = 0.020 \pm 0.001$
Clusters	$\Omega_B/\Omega_M = (0.07 \pm 0.007)h^{-3/2} \quad (\text{X ray})$
	$= (0.08 \pm 0.01)h^{-1} \quad (\text{S - Z})$
Hubble constant	$h = 0.72 \pm 0.07$

The shape of the power spectrum of matter inhomogeneity depends upon $\Omega_M h$ and Ω_B/Ω_M , as well as other parameters. Currently, the best determination of these two parameters come from an analysis of 160,000 redshifts in the 2-degree Field Galaxy Redshift Survey (2dFGRS; Percival et al, 2001). The baryon-to-total matter ratio is determined from the presence of “baryon bumps” in the power spectrum; it is hardly a significant result at the moment and gives almost no weight to the determination of Ω_M and Ω_B . Early results from the Sloan Digital Sky Survey (SDSS) are consistent with the 2dFGRS numbers, $\Omega_M h = 0.19 \pm 0.04$ (Dodelson et al., 2002; Szalay et al., 2001).

Significant improvement in both $\Omega_M h$ and Ω_B/Ω_M can be expected when the 2dF and SDSS surveys have amassed and analyzed their full samples of 250,000 and 600,000 redshifts respectively. The results will also be more robust. For example, current analyses assume scale-invariant perturbations ($n = 1$) and there are still correlations between σ_8 and $\Omega_M h$. While cosmic-shear measurements of large-scale structure are not yet good enough to have significant leverage on Ω_M and Ω_B , they will be in the future (see e.g., van Waerbeke et al., 2001; Maoli et al., 2001).

The structure of the acoustic peaks in the CMB angular power spectrum depends upon the total matter density and the baryon density (in physical units): the ratio of the odd to even peaks pins down the baryon density and the height of the first peak is sensitive to the total matter density (see e.g., Hu et al, 1997). I have used the values extracted from the first-year DASI results (Pryke et al, 2002). (The correlations between the two densities are small.) A re-analysis of the 1998 flight of Boomerang yields similar values (Netterfield et al, 2002). Combining the two results – which I have not done – would reduce the error bars by about $\sqrt{2}$.

While the consistency of DASI, a radio interferometer operating at around 30 GHz, and Boomerang, bolometers in bands above 100 GHz, is very reassuring, there are still significant issues. Both groups used a limited number of parameters (7) and priors on the Hubble constant ($h > 0.45$) and optical depth to last scattering ($\tau < 0.4$). The parameters preclude a large contribution by tensor (gravitational wave) perturbations; while unlikely, relaxing that assumption can significantly change the inferred baryon density (see e.g., Wang, Tegmark and Zaldarriaga, 2002). With MAP and Planck on the horizon, significant improvement, both in the precision and the robustness of the results, lies ahead.

At present, the most precise determination of the physical baryon density comes from combining measurements of the primeval abundance of deuterium (see e.g., O’Meara

et al, 2001) with accurate theoretical predictions of the light-element abundances (see e.g., Burles et al, 2001). I follow the analysis of Burles et al (2001) in adopting $\Omega_B h^2 = 0.02 \pm 0.001$. Here too, there are still issues to be resolved: possible unidentified systematics in the determination of the primeval deuterium abundance; reliable values for the primeval ^4He and ^7Li abundances to compare with the values predicted from the deuterium-determined baryon density. Over the next decade, as more deuterium systems are found and the ^4He and ^7Li abundances are better understood, the accuracy and reliability of the BBN baryon density should improve.

The baryon-to-matter ratio in clusters can be determined by x-ray measurements alone and by measurements of the Sunyaev – Zel’dovich (SZ) distortion of the CMB combined with x-ray measurements. For x-ray measurements alone, I adopt the cluster baryon fraction determined from a sample of 45 clusters by Mohr et al (1998), and for the SZ/X-ray determination I use the cluster sample of Grego et al (2001).

Potential sources of systematic error remain. There is some clumping of cluster gas, which could lead to an over-estimation of the amount of gas. The baryon-to-total mass ratio does vary with radius, and the calibrations are done by comparison with numerical simulations. The treatment of gas dynamics in these simulations still involves approximations and assumptions. Better simulations, more observations of clusters with the SZ technique and x-ray should address the key issues and improve the reliability of this sampling technique.

The sensitivity of the abundance of clusters, both as a function of cluster mass and redshift, to the matter density has been used to estimate Ω_M (see e.g., Bahcall & Fan, 1998; Blanchard et al, 2000; Henry, 2000). However, the range in inferred values is broad, $\Omega_M = 0.1 - 1$, largely because of the exponentially important, but uncertain relation between cluster mass and x-ray temperature. For this reason, I do not include these measurements in my determination of the matter density.

Finally, the value of the Hubble constant is important in converting physical densities to fractions of critical density; I adopt the value determined by the Hubble Key Project: $h = 0.72 \pm 0.07$ (Freedman et al, 2001). The error is almost entirely due to systematics (the statistical error is only $\sigma_h = \pm 0.02$). The largest part of the systematic error budget is in the distance to the Large Magellanic Cloud. Possible dependence of the Cepheid period – luminosity relation on metallicity may also be an issue.

3. THE CASE FOR $\Omega_m = 0.33 \pm 0.035$

Having adopted these physical measurements as input data, it is straightforward to deduce the fractions of critical density contributed by matter and by baryons, as well as the Hubble constant. But first, consider the consistency of these measurements. The physical baryon density ($\Omega_B h^2$) is determined by BBN, CMB anisotropy and the power spectrum + Hubble constant:

$$\begin{aligned} \Omega_B h^2 &= 0.020 \pm 0.001 && \text{BBN} \\ &= 0.022^{+0.004}_{-0.003} && \text{CMB} \\ &= 0.022 \pm 0.011 && \text{Power Spectrum} + H_0 \end{aligned} \quad (6)$$

These three, independent determinations of the baryon density are clearly consistent, giving one confidence in the

case for a low baryon density ($\rho_B \approx 4 \times 10^{-31} \text{ g cm}^{-3}$). They involve very different physics – nuclear reactions when the Universe was seconds old, gravity-driven acoustic oscillations when the Universe was around 400,000 yrs old, and the inhomogeneity in the distribution of matter in the Universe today – and thus also provide an important test the consistency of the big-bang framework and general relativity.

Next, consider the ratio of the total matter density to the baryon density:

$$\begin{aligned} \Omega_M/\Omega_B &= 7.2 \pm 2.1 && \text{CMB} \\ &= 6.7 \pm 3.1 && \text{Power Spectrum} \\ &= 9.0 \pm 1.4 && \text{Clusters (SZ) + } H_0 \\ &= 8.7 \pm 1.6 && \text{Clusters (x-ray) + } H_0 \end{aligned}$$

Again, all four measurements are clearly consistent and involve different physics – gravity driven acoustic oscillations, inhomogeneity in the distribution of matter today, and cluster dynamics.

Using standard techniques (and flat priors) I have constructed the likelihood function for Ω_M , Ω_B and h . From this, a posteriori probability distributions and credible ranges are calculated by marginalizing over one or two of these quantities in the usual way. The $1 - \sigma$ ranges are:

$$\begin{aligned} \Omega_B &= 0.039 \pm 0.0075 && \Omega_M = 0.33 \pm 0.035 \\ h &= 0.69 \pm 0.06 \end{aligned} \quad (7)$$

While the one-dimensional probability distributions are not perfectly Gaussian, the 68% and 95% credible ranges match pretty well with these 1σ error flags.

The value for the baryon fraction is largely driven by BBN and H_0 ; alone they imply $\Omega_B = 0.0385 \pm 0.0077$. Likewise, the Hubble constant is largely driven by its direct determination. The different scalings of the baryon-to-matter ratios with Hubble constant also provide important leverage, which can be seen in two ways: (1) the joint determination of h has a slightly smaller uncertainty than the direct measurement alone, ± 0.06 vs. ± 0.07 ; (2) if one arbitrarily doubles the uncertainty in the Hubble constant, $h = 0.72 \pm 0.14$, and carries out the same analysis, the results do not change dramatically:

$$\begin{aligned} \Omega_B &= 0.040 \pm 0.012 && \Omega_M = 0.33 \pm 0.045 \\ h &= 0.69 \pm 0.075 \end{aligned} \quad (8)$$

To investigate the robustness of my estimates for the matter and baryon densities I have studied their sensitivity to the individual input data. To investigate sensitivity to scale-dependent bias on $\Omega_M h$ I increased $\sigma_{\Omega_M h}$ by a factor of 4. The central values for Ω_M and Ω_B were unchanged and their errors increased to ± 0.044 and ± 0.0076 respectively. One by one I doubled the errors for the other input data; the central values for Ω_M and Ω_B changed little and the errors grew to at most ± 0.05 and ± 0.008 respectively. One minor trend was observed: enlarging the error flags on both the X-ray and SZ cluster baryon fractions by a factor of 4, to significantly reduce the weight given to the cluster baryon fraction, decreased the estimate for the matter density by about 1σ , $\Omega_M = 0.29 \pm 0.05$.

4. CONCLUDING REMARKS

Based upon present measurements of physical quantities not tied to the distribution of light I conclude that: $\Omega_M = 0.33 \pm 0.035$ and $\Omega_B = 0.039 \pm 0.0075$. While potential sources of systematic error remain, I have shown that these mass-density determinations are robust, and do not depend strongly upon any one measurement.

The precision and reliability of these determinations should improve over the next decade. With Planck and MAP the uncertainty in both $\Omega_M h^2$ and $\Omega_B h^2$ is likely to drop to the percent level (see e.g., Eisenstein et al, 1999); the cluster and power spectrum derived quantities are likely to improve by at least a factor of two. The situation with the Hubble constant is more difficult to predict.

To illustrate the potential for improvement, I have reduced the uncertainty in $\Omega_M h^2$ and $\Omega_B h^2$ to 1%, in $\Omega_M h$ to 5%, in Ω_B/Ω_M to 20%, and in the cluster baryon fractions and Hubble constant by a factor of two. The projected 1σ uncertainties for Ω_M and Ω_B drop to ± 0.012 and ± 0.0017 respectively.

Already the physically based matter density has interesting implications. First, combining it with the CMB determination of the total mass/energy density, $\Omega_0 = 1 \pm 0.05$ (Hanany et al, 2000; Netterfield et al, 2002; Pryke et al, 2002), makes a very strong case for an additional component to the Universe (referred to as dark energy) that is smoothly distributed with $\Omega_X = 0.67 \pm 0.06$. This evidence for the dark energy bolsters significantly the direct evidence for dark energy from supernovae (Perlmutter et al, 1999; Riess et al, 1998).

Second, the physically based matter density is about a factor of 2 larger than that determined from the mass-to-light ratios of clusters, $\Omega_M = 0.19 \pm 0.04$ (Carlberg et al, 1997) and 0.16 ± 0.05 (Bahcall et al, 2000). This suggests that baryons in the cluster environment produce twice as much light as in the field. Going a step further, this implies the fraction of baryons that become stars is about a factor of two higher. The SDSS will determine mass-to-light ratio for a stacked sample of millions of field galaxies in five-color bands using weak lensing and will test this hypothesis by providing a reliable estimate of $\langle M/L \rangle$.

Finally, in probing the nature of the dark energy, independent knowledge of the matter density is crucial for breaking the degeneracy between the dark energy equation-of-state ($w_X \equiv p_X/\rho_X$) and Ω_M (see e.g., Maor et al, 2001; Weller & Albrecht, 2001; Huterer & Turner, 2001). The value deduced here is essentially independent of the nature of the dark energy (assuming a flat Universe), and the current one-sigma uncertainty would already help significantly to break the degeneracy in future determinations of w_X .

Acknowledgments. This work was supported by the DoE (at Chicago and Fermilab), the NSF (through the CfCP) and by the NASA (at Fermilab by grant NAG 5-7092). I thank the participants of the 2002 Workshop on Large-scale Structure at the Aspen Center for Physics for helpful comments.

REFERENCES

- Blanchard, A., Sadat, R., Bartlett, J. & Le Dour, M. 2000, in Clustering at High Redshift (ASP Conference Series, vol. 200), eds. A. Mazure, O. Le Fevre & V. Le Brun, p. 158
Bahcall, N., Lubin, L. & Dorman, V. 1995, ApJ, 447, L81
Bahcall, N. et al 2000, ApJ, 541, 1
Burles, S., Nollett, K.M. & Turner, M.S. 2001, ApJ, 552, 1
Carlberg, R. et al 1997, ApJ, 478, 462
Dekel, A. 1994, ARA&A, 32, 371
Dekel, A., Burstein, D. & White, S.D.M. 1997, in Critical Dialogues in Cosmology, ed. N. Turok (WSPC, Singapore), p. 175
Dodelson, S. et al (SDSS Collaboration) 2002, ApJ, 572, 140
Eisenstein, D., Hu, W. & Tegmark, M. 1999, ApJ, 518, 2
Faber, S. & Gallagher, J. 1979, ARA&A, 17, 135
Fan, X. & Bahcall, N. 1998, ApJ, 504, 1
Fischer, P. et al (SDSS) 2000, AJ, 120, 1198
Freedman, W. et al 2001, ApJ, 553, 47
Grego, L. et al 2001, ApJ, 552, 2
Hanany, S. et al 2000, ApJ, 545, L5
Henry, J.P. 2000, ApJ, 534, 565
Hu, W., Sugiyama, N., & Silk, J. 1997, Nature, 386, 37
Huterer, D. & Turner, M.S. 2001, Phys. Rev. D, 64, 123527
A. Maoli et al 2001, *ibid.*, 368, 766
Maor, I., Burstein, R. & Steinhardt, P.J. 2001, Phys. Rev. Lett., 86, 6; 87, 049901 (E)
Mohr, J. et al 1998, ApJ, 517, 627
Netterfield, C.B. et al 2002, ApJ, 571, 604
O'Meara, J.M. et al 2001, ApJ, 552, 718
Percival, W.J. et al 2001, MNRAS, 327, 129
Perlmutter, S. et al 1999, ApJ, 517, 565
Primack, J. 2001, in Sources and Detection of Dark Matter and Dark Energy in the Universe, ed. D. Cline (Springer-Verlag, Berlin), p. 3 (astro-ph/0007187)
Pryke, C. et al 2002, ApJ, 568, 46
Riess, A. et al 1998, AJ, 116, 1009
Szalay, A. et al (SDSS Collaboration) 2001, astro-ph/0107421
Turner, M.S. 2000, Physica Scripta, T85, 210
van Waerbeke, L. et al 2001, *ibid.*, 375, 757
Wang, X., Tegmark, M. & Zaldarriaga 2002, Phys. Rev. D, 65, 123001
Weller, J. & Albrecht, A. 2001, Phys. Rev. Lett., 86, 1939
White, S.D.M. et al 1993, Nature, 366, 429
Willick, J. & Strauss, M. 1998, ApJ, 507, 64